

## PROBLEMS


### Section 1.2 Units

### Section 1.3 The Role of Units in Problem Solving


- 1.** **GO** A student sees a newspaper ad for an apartment that has 1330 square feet ( $\text{ft}^2$ ) of floor space. How many square meters of area are there?
- 2.** Bicyclists in the Tour de France reach speeds of 34.0 miles per hour (mi/h) on flat sections of the road. What is this speed in **(a)** kilometers per hour (km/h) and **(b)** meters per second (m/s)?
- 3. ssm** Vesna Vulovic survived the longest fall on record without a parachute when her plane exploded and she fell 6 miles, 551 yards. What is this distance in meters?
- 4.** Suppose a man's scalp hair grows at a rate of 0.35 mm per day. What is this growth rate in feet per century?
- 5.** Given the quantities  $a = 9.7 \text{ m}$ ,  $b = 4.2 \text{ s}$ ,  $c = 69 \text{ m/s}$ , what is the value of the quantity  $d = a^3/(cb^2)$ ?
- 6.** Consider the equation  $v = \frac{1}{3} zxt^2$ . The dimensions of the variables  $v$ ,  $x$ , and  $t$  are  $[\text{L}]/[\text{T}]$ ,  $[\text{L}]$ , and  $[\text{T}]$ , respectively. The numerical factor 3 is dimensionless. What must be the dimensions of the variable  $z$ , such that both sides of the equation have the same dimensions? Show how you determined your answer.
- 7. ssm** A bottle of wine known as a magnum contains a volume of 1.5 liters. A bottle known as a jeroboam contains 0.792 U.S. gallons. How many magnums are there in one jeroboam?
- 8.** The CGS unit for measuring the viscosity of a liquid is the poise (P):  $1 \text{ P} = 1 \text{ g}/(\text{s} \cdot \text{cm})$ . The SI unit for viscosity is the  $\text{kg}/(\text{s} \cdot \text{m})$ . The viscosity of water at  $0^\circ \text{C}$  is  $1.78 \times 10^{-3} \text{ kg}/(\text{s} \cdot \text{m})$ . Express this viscosity in poise.
- 9. GO** Azelastine hydrochloride is an antihistamine nasal spray. A standard-size container holds one fluid ounce (oz) of the liquid. You are searching for this medication in a European drugstore and are asked how many milliliters (mL) there are in one fluid ounce. Using the following conversion factors, determine the number of milliliters in a volume of one fluid ounce:  $1 \text{ gallon (gal)} = 128 \text{ oz}$ ,  $3.785 \times 10^{-3} \text{ cubic meters (m}^3\text{)} = 1 \text{ gal}$ , and  $1 \text{ mL} = 10^{-6} \text{ m}^3$ .
- \*10. GO** A partly full paint can has 0.67 U.S. gallons of paint left in it. **(a)** What is the volume of the paint in cubic meters? **(b)** If all the remaining paint is used to coat a wall evenly (wall area =  $13 \text{ m}^2$ ), how thick is the layer of wet paint? Give your answer in meters.
- \*11. ssm** A spring is hanging down from the ceiling, and an object of mass  $m$  is attached to the free end. The object is pulled down, thereby stretching the spring, and then released. The object oscillates up and down, and the time  $T$  required for one complete up-and-down oscillation is given by the equation  $T = 2\pi\sqrt{m/k}$ , where  $k$  is known as the spring constant. What must be the dimension of  $k$  for this equation to be dimensionally correct?

### Section 1.4 Trigonometry

- 12.** You are driving into St. Louis, Missouri, and in the distance you see the famous Gateway to the West arch. This monument rises to a height of 192 m. You estimate your line of sight with the top of the arch to be  $2.0^\circ$  above the horizontal. Approximately how far (in kilometers) are you from the base of the arch?
- 13. ssm** A highway is to be built between two towns, one of which lies 35.0 km south and 72.0 km west of the other. What is the shortest length of highway that can be built between the two towns, and at what angle would this highway be directed with respect to due west?
- 14.** A hill that has a 12.0% grade is one that rises 12.0 m vertically for every 100.0 m of distance in the horizontal direction. At what angle is such a hill inclined above the horizontal?
- 15.** The corners of a square lie on a circle of diameter  $D = 0.35 \text{ m}$ . Each side of the square has a length  $L$ . Find  $L$ .
- 16.** The drawing shows a person looking at a building on top of which an antenna is mounted. The horizontal distance between the person's eyes and the building is 85.0 m. In part *a* the person is looking at the base of the antenna, and his line of sight makes an angle of  $35.0^\circ$  with the horizontal. In part *b* the person is looking at the top of the antenna, and his line of sight makes an angle of  $38.0^\circ$  with the horizontal. How tall is the antenna?




17. The two hot-air balloons in the drawing are 48.2 and 61.0 m above the ground. A person in the left balloon observes that the right balloon is  $13.3^\circ$  above the horizontal. What is the horizontal distance  $x$  between the two balloons?




\*18. What is the value of each of the angles of a triangle whose sides are 95, 150, and 190 cm in length?  
(Hint: Consider using the law of cosines given in Appendix E.)

\*19. **mmh** The drawing shows sodium and chloride ions positioned at the corners of a cube that is part of the crystal structure of sodium chloride (common table salt). The edges of the cube are each 0.281 nm (1 nm = 1 nanometer =  $10^{-9}$  m) in length. What is the value of the angle  $\theta$  in the drawing?




\*20. **GO** A person is standing at the edge of the water and looking out at the ocean (see the drawing). The height of the person's eyes above the water is  $h = 1.6$  m, and the radius of the earth is  $R = 6.38 \times 10^6$  m. **(a)** How far is it to the horizon? In other words, what is the distance  $d$  from the person's eyes to the horizon? (Note: At the horizon the angle between the line of sight and the radius of the earth is  $90^\circ$ .) **(b)** Express this distance in miles.



\*21. **ssm** Three deer, A, B, and C, are grazing in a field. Deer B is located 62 m from deer A at an angle of  $51^\circ$  north of west. Deer C is located  $77^\circ$  north of east relative to deer A. The distance between deer B and C is 95 m. What is the distance between deer A and C? (Hint: Consider the law of cosines given in Appendix E.)

\*22. An aerialist on a high platform holds on to a trapeze attached to a support by an 8.0-m cord. (See the drawing.) Just before he jumps off the platform, the cord makes an angle of  $41^\circ$  with the vertical. He jumps,

swings down, then back up, releasing the trapeze at the instant it is 0.75 m below its initial height. Calculate the angle  $\theta$  that the trapeze cord makes with the vertical at this instant.



## Section 1.6 Vector Addition and Subtraction

**23. ssm** (a) Two workers are trying to move a heavy crate. One pushes on the crate with a force  $\vec{A}$ , which has a magnitude of 445 newtons and is directed due west. The other pushes with a force  $\vec{B}$ , which has a magnitude of 325 newtons and is directed due north. What are the magnitude and direction of the resultant force  $\vec{A} + \vec{B}$  applied to the crate? (b) Suppose that the second worker applies a force  $-\vec{B}$  instead of  $\vec{B}$ . What then are the magnitude and direction of the resultant force  $\vec{A} - \vec{B}$  applied to the crate? In both cases express the direction relative to due west.

**24.** A force vector  $\vec{F}_1$  points due east and has a magnitude of 200 newtons. A second force  $\vec{F}_2$  is added to  $\vec{F}_1$ . The resultant of the two vectors has a magnitude of 400 newtons and points along the east/west line. Find the magnitude and direction of  $\vec{F}_2$ . Note that there are two answers.

**25. ssm** Consider the following four force vectors:

$\vec{F}_1 = 50.0$  newtons, due east

$\vec{F}_2 = 10.0$  newtons, due east

$\vec{F}_3 = 40.0$  newtons, due west


$\vec{F}_4 = 30.0$  newtons, due west

Which two vectors add together to give a resultant with the smallest magnitude, and which two vectors add to give a resultant with the largest magnitude? In each case specify the magnitude and direction of the resultant.


**26. GO** Vector  $\vec{A}$  has a magnitude of 63 units and points due west, while vector  $\vec{B}$  has the same magnitude and points due south. Find the magnitude and direction of (a)  $\vec{A} + \vec{B}$  and (b)  $\vec{A} - \vec{B}$ . Specify the directions relative to due west.

**27.** Two bicyclists, starting at the same place, are riding toward the same campground by two different routes. One cyclist rides 1080 m due east and then turns due north and travels another 1430 m before reaching the campground. The second cyclist starts out by heading due north for 1950 m and then turns and heads directly toward the campground. (a) At the turning point, how far is the second cyclist from the campground? (b) In what direction (measured relative to due east) must the second cyclist head during the last part of the trip?

**28. GO** The drawing shows a triple jump on a checkerboard, starting at the center of square A and ending on the center of square B. Each side of a square measures 4.0 cm. What is the magnitude of the displacement of the colored checker during the triple jump?




29. Given the vectors  $\vec{P}$  and  $\vec{Q}$  shown on the grid, sketch and calculate the magnitudes of the vectors (a)  $\vec{M} = \vec{P} + \vec{Q}$  and (b)  $\vec{K} = 2\vec{P} - \vec{Q}$ . Use the tail-to-head method and express the magnitudes in centimeters with the aid of the grid scale shown in the drawing.




\*30. **mmh** Vector  $\vec{A}$  has a magnitude of 12.3 units and points due west. Vector  $\vec{B}$  points due north. (a) What is the magnitude of  $\vec{B}$  if  $\vec{A} + \vec{B}$  has a magnitude of 15.0 units? (b) What is the direction of  $\vec{A} + \vec{B}$  relative to due west? (c) What is the magnitude of  $\vec{B}$  if  $\vec{A} - \vec{B}$  has a magnitude of 15.0 units? (d) What is the direction of  $\vec{A} - \vec{B}$  relative to due west?

\*31. **ssm** A car is being pulled out of the mud by two forces that are applied by the two ropes shown in the drawing. The dashed line in the drawing bisects the  $30.0^\circ$  angle. The magnitude of the force applied by each rope is 2900 newtons. Arrange the force vectors tail to head and use the graphical technique to answer the following questions. (a) How much force would a single rope need to apply to accomplish the same effect as the two forces added together? (b) How would the single rope be directed relative to the dashed line?



\*32. **GO** A jogger travels a route that has two parts. The first is a displacement  $\vec{A}$  of 2.50 km due south, and the second involves a displacement  $\vec{B}$  that points due east. (a) The resultant displacement  $\vec{A} + \vec{B}$  has a magnitude of 3.75 km. What is the magnitude of  $\vec{B}$ , and what is the direction of  $\vec{A} + \vec{B}$  relative to due south? (b) Suppose that  $\vec{A} - \vec{B}$  had a magnitude of 3.75 km. What then would be the magnitude of  $\vec{B}$ , and what is the direction of  $\vec{A} - \vec{B}$  relative to due south?

\*33. At a picnic, there is a contest in which hoses are used to shoot water at a beach ball from three directions. As a result, three forces act on the ball,  $\vec{F}_1$ ,  $\vec{F}_2$ , and  $\vec{F}_3$  (see the drawing). The magnitudes of  $\vec{F}_1$  and  $\vec{F}_2$  are  $F_1 = 50.0$  newtons and  $F_2 = 90.0$  newtons. Using a scale drawing and the graphical technique, determine (a) the magnitude of  $\vec{F}_3$  and (b) the angle  $\theta$  such that the resultant force acting on the ball is zero.




### Section 1.7 The Components of a Vector

34. **GO** A force vector has a magnitude of 575 newtons and points at an angle of  $36.0^\circ$  below the positive  $x$  axis. What are (a) the  $x$  scalar component and (b) the  $y$  scalar component of the vector?

35. **ssm** Vector  $\vec{A}$  points along the  $+y$  axis and has a magnitude of 100.0 units. Vector  $\vec{B}$  points at an angle of  $60.0^\circ$  above the  $+x$  axis and has a magnitude of 200.0 units. Vector  $\vec{C}$  points along the  $+x$  axis and has a magnitude of 150.0 units. Which vector has (a) the largest  $x$  component and (b) the largest  $y$  component?

36. Soccer player #1 is 8.6 m from the goal (see the drawing). If she kicks the ball directly into the net, the ball has a displacement labeled  $\vec{A}$ . If, on the other hand, she first kicks it to player #2, who then kicks it into the net, the ball undergoes two successive displacements,  $\vec{A}_y$  and  $\vec{A}_x$ . What are the magnitudes and directions of  $\vec{A}_x$  and  $\vec{A}_y$ ?




Problem 36

37. The components of vector  $\vec{A}$  are  $A_x$  and  $A_y$  (both positive), and the angle that it makes with respect to the positive  $x$  axis is  $\theta$ . Find the angle  $\theta$  if the components of the displacement vector  $\vec{A}$  are (a)  $A_x = 12$  m and  $A_y = 12$  m, (b)  $A_x = 17$  m and  $A_y = 12$  m, and (c)  $A_x = 12$  m and  $A_y = 17$  m.
38. During takeoff, an airplane climbs with a speed of 180 m/s at an angle of  $34^\circ$  above the horizontal. The speed and direction of the airplane constitute a vector quantity known as the velocity. The sun is shining directly overhead. How fast is the shadow of the plane moving along the ground? (That is, what is the magnitude of the horizontal component of the plane's velocity?)
39. **ssm** The  $x$  vector component of a displacement vector  $\vec{r}$  has a magnitude of 125 m and points along the negative  $x$  axis. The  $y$  vector component has a magnitude of 184 m and points along the negative  $y$  axis. Find the magnitude and direction of  $\vec{r}$ . Specify the direction with respect to the negative  $x$  axis.
40. Your friend has slipped and fallen. To help her up, you pull with a force  $\vec{F}$ , as the drawing shows. The vertical component of this force is 130 newtons, and the horizontal component is 150 newtons. Find (a) the magnitude of  $\vec{F}$  and (b) the angle  $\theta$ .




41. The displacement vector  $\vec{A}$  has scalar components of  $A_x = 80.0$  m and  $A_y = 60.0$  m. The displacement vector  $\vec{B}$  has a scalar component of  $B_x = 60.0$  m and a magnitude of  $B = 75.0$  m. The displacement vector  $\vec{C}$  has a magnitude of  $C = 100.0$  m and is directed at an angle of  $36.9^\circ$  above the  $+x$  axis. Two of these vectors are equal. Determine which two, and support your choice with a calculation.
- \*42. Two racing boats set out from the same dock and speed away at the same constant speed of 101 km/h for half an hour (0.500 h), the blue boat headed  $25.0^\circ$  south of west, and the green boat headed  $37.0^\circ$  south of west. During this half hour (a) how much farther west does the blue boat travel, compared to the green boat, and (b) how much farther south does the green boat travel, compared to the blue boat? Express your answers in km.
- \*43. **ssm mmh** The magnitude of the force vector  $\vec{F}$  is 82.3 newtons. The  $x$  component of this vector is directed along the  $+x$  axis and has a magnitude of 74.6 newtons. The  $y$  component points along the  $+y$  axis. (a) Find the direction of  $\vec{F}$  relative to the  $+x$  axis. (b) Find the component of  $\vec{F}$  along the  $+y$  axis.
- \*\*44. The drawing shows a force vector that has a magnitude of 475 newtons. Find the (a)  $x$ , (b)  $y$ , and (c)  $z$  components of the vector.



## Section 1.8 Addition of Vectors by Means of Components

**45. ssm** Consult Multiple-Concept Example 9 in preparation for this problem. A golfer, putting on a green, requires three strokes to “hole the ball.” During the first putt, the ball rolls 5.0 m due east. For the second putt, the ball travels 2.1 m at an angle of  $20.0^\circ$  north of east. The third putt is 0.50 m due north. What displacement (magnitude and direction relative to due east) would have been needed to “hole the ball” on the very first putt?


**46. P11** The three displacement vectors in the drawing have magnitudes of  $A = 5.00$  m,  $B = 5.00$  m, and  $C = 4.00$  m. Find the resultant (magnitude and directional angle) of the three vectors by means of the component method. Express the directional angle as an angle above the positive or negative  $x$  axis.



Problem 46

**47. mmh** Multiple-Concept Example 9 reviews the concepts that play a role in this problem. Two forces are applied to a tree stump to pull it out of the ground. Force  $\vec{F}_A$  has a magnitude of 2240 newtons and points  $34.08^\circ$  south of east, while force  $\vec{F}_B$  has a magnitude of 3160 newtons and points due south. Using the component method, find the magnitude and direction of the resultant force  $\vec{F}_A + \vec{F}_B$  that is applied to the stump. Specify the direction with respect to due east.


**48. P10** A baby elephant is stuck in a mud hole. To help pull it out, game keepers use a rope to apply a force  $\vec{F}_A$ , as part *a* of the drawing shows. By itself, however, force  $\vec{F}_A$  is insufficient. Therefore, two additional forces  $\vec{F}_B$  and  $\vec{F}_C$  are applied, as in part *b* of the drawing. Each of these additional forces has the same magnitude  $F$ . The magnitude of the resultant force acting on the elephant in part *b* of the drawing is  $k$  times larger than that in part *a*. Find the ratio  $F/F_A$  when  $k = 2.00$ .



Problem 48

**49.** Displacement vector  $\vec{A}$  points due east and has a magnitude of 2.00 km. Displacement vector  $\vec{B}$  points due north and has a magnitude of 3.75 km. Displacement vector  $\vec{C}$  points due west and has a magnitude of 2.50 km. Displacement vector  $\vec{D}$  points due south and has a magnitude of 3.00 km. Find the magnitude and direction (relative to due west) of the resultant vector  $\vec{A} + \vec{B} + \vec{C} + \vec{D}$ .

**50.** Multiple-Concept Example 9 provides background pertinent to this problem. The magnitudes of the four displacement vectors shown in the drawing are  $A = 16.0$  m,  $B = 11.0$  m,  $C = 12.0$  m, and  $D = 26.0$  m. Determine the magnitude and directional angle for the resultant that occurs when these vectors are added together.




**51. mmh** On a safari, a team of naturalists sets out toward a research station located 4.8 km away in a direction  $42^\circ$  north of east. After traveling in a straight line for 2.4 km, they stop and discover that they have been traveling  $22^\circ$  north of east, because their guide misread his compass. What are (a) the magnitude and

(b) the direction (relative to due east) of the displacement vector now required to bring the team to the research station?

\*52.  Two geological field teams are working in a remote area. A global positioning system (GPS) tracker at their base camp shows the location of the first team as 38 km away,  $19^\circ$  north of west, and the second team as 29 km away,  $35^\circ$  east of north. When the first team uses its GPS to check the position of the second team, what does the GPS give for the second team's (a) distance from them and (b) direction, measured from due east?

\*53. **ssm** A sailboat race course consists of four legs, defined by the displacement vectors  $\vec{A}$ ,  $\vec{B}$ ,  $\vec{C}$ , and  $\vec{D}$ , as the drawing indicates. The magnitudes of the first three vectors are  $A = 3.20\text{ km}$ ,  $B = 5.10\text{ km}$ , and  $C = 4.80\text{ km}$ . The finish line of the course coincides with the starting line. Using the data in the drawing, find the distance of the fourth leg and the angle  $\theta$ .



\*54. Multiple-Concept Example 9 deals with the concepts that are important in this problem. A grasshopper makes four jumps. The displacement vectors are (1) 27.0 cm, due west; (2) 23.0 cm,  $35.0^\circ$  south of west; (3) 28.0 cm,  $55.0^\circ$  south of east; and (4) 35.0 cm,  $63.0^\circ$  north of east. Find the magnitude and direction of the resultant displacement. Express the direction with respect to due west.

\*55. **mmh** Vector  $\vec{A}$  has a magnitude of 145 units and points  $35.0^\circ$  north of west. Vector  $\vec{B}$  points  $65.0^\circ$  east of north. Vector  $\vec{C}$  points  $15.0^\circ$  west of south. These three vectors add to give a resultant vector that is zero. Using components, find the magnitudes of (a) vector  $\vec{B}$  and (b) vector  $\vec{C}$ .

\*56. The route followed by a hiker consists of three displacement vectors  $\vec{A}$ ,  $\vec{B}$ , and  $\vec{C}$ . Vector  $\vec{A}$  is along a measured trail and is 1550 m in a direction  $25.0^\circ$  north of east. Vector  $\vec{B}$  is not along a measured trail, but the hiker uses a compass and knows that the direction is  $41.0^\circ$  east of south. Similarly, the direction of vector  $\vec{C}$  is  $35.0^\circ$  north of west. The hiker ends up back where she started. Therefore, it follows that the resultant displacement is zero, or  $\vec{A} + \vec{B} + \vec{C} = 0$ . Find the magnitudes of (a) vector  $\vec{B}$  and (b) vector  $\vec{C}$ .

## ADDITIONAL PROBLEMS

57. A chimpanzee sitting against his favorite tree gets up and walks 51 m due east and 39 m due south to reach a termite mound, where he eats lunch. (a) What is the shortest distance between the tree and the termite mound? (b) What angle does the shortest distance make with respect to due east?

58. A monkey is chained to a stake in the ground. The stake is 3.00 m from a vertical pole, and the chain is 3.40 m long. How high can the monkey climb up the pole?


59. **ssm** The speed of an object and the direction in which it moves constitute a vector quantity known as the velocity. An ostrich is running at a speed of 17.0 m/s in a direction of  $68.0^\circ$  north of west. What is the magnitude of the ostrich's velocity component that is directed (a) due north and (b) due west?

60.  The volume of liquid flowing per second is called the volume flow rate  $Q$  and has the dimensions of  $[L]^3/[T]$ . The flow rate of a liquid through a hypodermic needle during an injection can be estimated with the following equation:

$$Q = \frac{\pi R^n (P_2 - P_1)}{8\eta L}$$

The length and radius of the needle are  $L$  and  $R$ , respectively, both of which have the dimension  $[L]$ . The pressures at opposite ends of the needle are  $P_2$  and  $P_1$ , both of which have the dimensions of  $[M]/\{[L][T]^2\}$ . The symbol  $\eta$  represents the viscosity of the liquid and has the dimensions of  $[M]/\{[L][T]\}$ . The symbol  $\pi$  stands for pi and, like the number 8 and the exponent  $n$ , has no dimensions. Using dimensional analysis, determine the value of  $n$  in the expression for  $Q$ .

61. An ocean liner leaves New York City and travels  $18.0^\circ$  north of east for 155 km. How far east and how far north has it gone? In other words, what are the magnitudes of the components of the ship's displacement vector in the directions (a) due east and (b) due north?
62.  A pilot flies her route in two straight-line segments. The displacement vector  $\vec{A}$  for the first segment has a magnitude of 244 km and a direction  $30.0^\circ$  north of east. The displacement vector  $\vec{B}$  for the second segment has a magnitude of 175 km and a direction due west. The resultant displacement vector is  $\vec{R} = \vec{A} + \vec{B}$  and makes an angle  $\theta$  with the direction due east. Using the component method, find the magnitude of  $\vec{R}$  and the directional angle  $\theta$ .
63. **ssm** A circus performer begins his act by walking out along a nearly horizontal high wire. He slips and falls to the safety net, 25.0 ft below. The magnitude of his displacement from the beginning of the walk to the net is 26.7 ft. (a) How far out along the high wire did he walk? (b) Find the angle that his displacement vector makes below the horizontal.
- \*64. A force vector points at an angle of  $52^\circ$  above the  $+x$  axis. It has a  $y$  component of +290 newtons. Find (a) the magnitude and (b) the  $x$  component of the force vector.
- \*65. **ssm** Vector  $\vec{A}$  has a magnitude of 6.00 units and points due east. Vector  $\vec{B}$  points due north. (a) What is the magnitude of  $\vec{B}$ , if the vector  $\vec{A} + \vec{B}$  points  $60.0^\circ$  north of east? (b) Find the magnitude of  $\vec{A} + \vec{B}$ .
- \*66.  Three forces act on an object, as indicated in the drawing. Force  $\vec{F}_1$  has a magnitude of 21.0 newtons (21.0 N) and is directed  $30.0^\circ$  to the left of the  $+y$  axis. Force  $\vec{F}_2$  has a magnitude of 15.0 N and points along the  $+x$  axis. What must be the magnitude and direction (specified by the angle  $\theta$  in the drawing) of the third force  $\vec{F}_3$  such that the vector sum of the three forces is 0 N?



- \*67. Before starting this problem, review Conceptual Example 7. The force vector  $\vec{F}_A$  has a magnitude of 90.0 newtons and points due east. The force vector  $\vec{F}_B$  has a magnitude of 135 newtons and points  $75^\circ$  north of east. Use the graphical method and find the magnitude and direction of (a)  $\vec{F}_A - \vec{F}_B$  (give the direction with respect to due east) and (b)  $\vec{F}_B - \vec{F}_A$  (give the direction with respect to due west).
- \*68.  You live in the building on the left in the drawing, and a friend lives in the other building. The two of you are having a discussion about the heights of the buildings, and your friend claims that the height of his building is more than 1.50 times the height of yours. To resolve the issue you climb to the roof of your building and estimate that your line of sight to the top edge of the other building makes an angle of  $21^\circ$  above the horizontal, whereas your line of sight to the base of the other building makes an angle of  $52^\circ$  below the horizontal. Determine the ratio of the height of the taller building to the height of the shorter building. State whether your friend is right or wrong.



- \*\*69. What are the  $x$  and  $y$  components of the vector that must be added to the following three vectors, so that the sum of the four vectors is zero? Due east is the  $+x$  direction, and due north is the  $+y$  direction.

$\vec{A} = 113$  units,  $60.0^\circ$  south of west


$\vec{B} = 222$  units,  $35.0^\circ$  south of east

$\vec{C} = 177$  units,  $23.0^\circ$  north of east


## CONCEPTS AND CALCULATIONS PROBLEMS

This chapter has presented an introduction to the mathematics of trigonometry and vectors, which will be used throughout the text. In this section we apply some of the important features of this mathematics, and review some concepts that can help in anticipating some of the characteristics of the numerical answers.

**\*70.**  The figure shows two displacement vectors  $\vec{A}$  and  $\vec{B}$ . Vector  $\vec{A}$  points at an angle of  $22^\circ$  above the  $x$  axis and has an unknown magnitude. Vector  $\vec{B}$  has an  $x$  component  $B_x = 35.0$  m and has an unknown  $y$  component  $B_y$ . These two vectors are equal. *Concepts:* (i) What does the condition that vector  $\vec{A}$  equals  $\vec{B}$  imply about the magnitudes and directions of the vectors? (ii) What does the condition that vector  $\vec{A}$  equals  $\vec{B}$  imply about the  $x$  and  $y$  components of the vectors? *Calculations:* Find the magnitude of  $\vec{A}$  and the value of  $B_y$ .




(a)



(b)

Problem 70

**\*71.**  **ssm** The figure shows three displacement vectors  $\vec{A}$ ,  $\vec{B}$ , and  $\vec{C}$ . These vectors are arranged in tail-to-head fashion, and they add together to give a resultant displacement  $\vec{R}$ , which lies along the  $x$  axis. Note that the vector  $\vec{B}$  is parallel to the  $x$  axis. *Concepts:* (i) How is the magnitude of  $\vec{A}$  related to its scalar components  $A_x$  and  $A_y$ ? (ii) Do any of the vectors in the figure have a zero value for either their  $x$  or  $y$  components? (If so, which ones?) (iii) What does the fact that  $\vec{A}$ ,  $\vec{B}$ , and  $\vec{C}$  add together to give  $\vec{R}$  tell you about the components of these vectors? *Calculations:* What is the magnitude of the vector  $\vec{A}$  and its directional angle  $\theta$ ?



Problem 71