



2019 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

3. A student is given 50.0 mL of a solution of Na_2CO_3 of unknown concentration. To determine the concentration of the solution, the student mixes the solution with excess 1.0 M $\text{Ca}(\text{NO}_3)_2(aq)$, causing a precipitate to form. The balanced equation for the reaction is shown below.



- (a) Write the net ionic equation for the reaction that occurs when the solutions of Na_2CO_3 and $\text{Ca}(\text{NO}_3)_2$ are mixed.
- (b) The diagram below is incomplete. Draw in the species needed to accurately represent the major ionic species remaining in the solution after the reaction has been completed.




The student filters and dries the precipitate of CaCO_3 (molar mass 100.1 g/mol) and records the data in the table below.

Volume of Na_2CO_3 solution	50.0 mL
Volume of 1.0 M $\text{Ca}(\text{NO}_3)_2$ added	100.0 mL
Mass of CaCO_3 precipitate collected	0.93 g

- (c) Determine the number of moles of Na_2CO_3 in the original 50.0 mL of solution.
- (d) The student realizes that the precipitate was not completely dried and claims that as a result, the calculated Na_2CO_3 molarity is too low. Do you agree with the student's claim? Justify your answer.
- (e) After the precipitate forms and is filtered, the liquid that passed through the filter is tested to see if it can conduct electricity. What would be observed? Justify your answer.

2019 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

The student decides to determine the molarity of the same Na_2CO_3 solution using a second method. When Na_2CO_3 is dissolved in water, $\text{CO}_3^{2-}(aq)$ hydrolyzes to form $\text{HCO}_3^-(aq)$, as shown by the following equation.



- (f) The student decides to first determine $[\text{OH}^-]$ in the solution, then use that result to calculate the initial concentration of $\text{CO}_3^{2-}(aq)$.
- Identify a laboratory method (not titration) that the student could use to collect data to determine $[\text{OH}^-]$ in the solution.
 - Explain how the student could use the measured value in part (f)(i) to calculate the initial concentration of $\text{CO}_3^{2-}(aq)$. (Do not do any numerical calculations.)
- (g) In the original Na_2CO_3 solution at equilibrium, is the concentration of $\text{HCO}_3^-(aq)$ greater than, less than, or equal to the concentration of $\text{CO}_3^{2-}(aq)$? Justify your answer.
- (h) The student needs to make a $\text{CO}_3^{2-}/\text{HCO}_3^-$ buffer. Is the Na_2CO_3 solution suitable for making a buffer with a pH of 6? Explain why or why not.