



AP[®] CHEMISTRY
2016 SCORING GUIDELINES

Question 3



To determine the molar mass of an unknown metal, M , a student reacts iodine with an excess of the metal to form the water-soluble compound MI_2 , as represented by the equation above. The reaction proceeds until all of the I_2 is consumed. The $MI_2(aq)$ solution is quantitatively collected and heated to remove the water, and the product is dried and weighed to constant mass. The experimental steps are represented below, followed by a data table.



Data for Unknown Metal Lab	
Mass of beaker	125.457 g
Mass of beaker + metal M	126.549 g
Mass of beaker + metal M + I_2	127.570 g
Mass of MI_2 , first weighing	1.284 g
Mass of MI_2 , second weighing	1.284 g

AP[®] CHEMISTRY
2016 SCORING GUIDELINES

Question 3 (continued)

- (a) Given that the metal M is in excess, calculate the number of moles of I₂ that reacted.

$127.570 - 126.549 = 1.021 \text{ g I}_2$ $1.021 \text{ g I}_2 \times \frac{1 \text{ mol I}_2}{253.80 \text{ g I}_2} = 0.004023 \text{ mol I}_2$	1 point is earned for the number of moles.
--	--

- (b) Calculate the molar mass of the unknown metal M.

Number of moles of I ₂ = number of moles of M $1.284 \text{ g MI}_2 - 1.021 \text{ g I}_2 = 0.263 \text{ g M}$ $\text{Molar mass of M} = \frac{0.263 \text{ g M}}{0.004023 \text{ mol M}} = 65.4 \text{ g/mol}$	1 point is earned for the number of grams of M. 1 point is earned for the molar mass.
---	--

The student hypothesizes that the compound formed in the synthesis reaction is ionic.

- (c) Propose an experimental test the student could perform that could be used to support the hypothesis. Explain how the results of the test would support the hypothesis if the substance was ionic.

The student could dissolve the compound in water or melt the compound and see if the solution/melt conducts electricity. If the solution/melt conducts electricity, mobile ions capable of carrying charge must be present, thus the compound is likely to be ionic. OR The student could heat the compound until it melts or boils. If the melting/boiling point is very high, then the compound is likely to be ionic.	1 point is earned for an appropriate test. 1 point is earned for explaining how the results would support the hypothesis.
--	--

The student hypothesizes that Br₂ will react with metal M more vigorously than I₂ did because Br₂ is a liquid at room temperature.

**AP[®] CHEMISTRY
2016 SCORING GUIDELINES**

Question 3 (continued)

- (d) Explain why I_2 is a solid at room temperature whereas Br_2 is a liquid. Your explanation should clearly reference the types and relative strengths of the intermolecular forces present in each substance.

<p>Both Br_2 and I_2 molecules are nonpolar molecules, therefore the only possible intermolecular forces are London dispersion forces.</p> <p>The London dispersion forces are stronger in I_2 because it is larger in size with more electrons and/or a more polarizable electron cloud. The stronger London dispersion forces in I_2 result in a higher melting point, which makes I_2 a solid at room temperature.</p>	<p>1 point is earned for identifying the forces in each substance as London dispersion forces.</p> <p>1 point is earned for explaining why the forces are stronger in I_2 than in Br_2.</p>
--	---

While cleaning up after the experiment, the student wishes to dispose of the unused solid I_2 in a responsible manner. The student decides to convert the solid I_2 to $I^-(aq)$ anion. The student has access to three solutions, $H_2O_2(aq)$, $Na_2S_2O_3(aq)$, and $Na_2S_4O_6(aq)$, and the standard reduction table shown below.

Half-reaction	E° (V)
$S_4O_6^{2-}(aq) + 2 e^- \rightarrow 2 S_2O_3^{2-}(aq)$	0.08
$I_2(s) + 2 e^- \rightarrow 2 I^-(aq)$	0.54
$O_2(g) + 2 H^+(aq) + 2 e^- \rightarrow H_2O_2(aq)$	0.68

- (e) Which solution should the student add to $I_2(s)$ to reduce it to $I^-(aq)$? Circle your answer below. Justify your answer and include a calculation of E° for the overall reaction.

$H_2O_2(aq)$

$Na_2S_2O_3(aq)$

$Na_2S_4O_6(aq)$

<p>[$Na_2S_2O_3(aq)$ should be circled.]</p> <p>The reaction between $S_2O_3^{2-}(aq)$ and $I_2(s)$ will be thermodynamically favorable because E° for the reaction is positive ($E^\circ = 0.54 - 0.08 = +0.46$ V), from which it follows that ΔG° is negative because $\Delta G^\circ = -nFE^\circ$.</p>	<p>1 point is earned for the correct choice.</p> <p>1 point is earned for a correct justification.</p>
---	--

- (f) Write the balanced net-ionic equation for the reaction between I_2 and the solution you selected in part (e).

$I_2 + 2 S_2O_3^{2-} \rightarrow 2 I^- + S_4O_6^{2-}$	1 point is earned for the correct equation.
---	---