


AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 3


A student is given 50.0 mL of a solution of Na_2CO_3 of unknown concentration. To determine the concentration of the solution, the student mixes the solution with excess 1.0 M $\text{Ca}(\text{NO}_3)_2(\text{aq})$, causing a precipitate to form. The balanced equation for the reaction is shown below.



- (a) Write the net ionic equation for the reaction that occurs when the solutions of Na_2CO_3 and $\text{Ca}(\text{NO}_3)_2$ are mixed.

$\text{Ca}^{2+}(\text{aq}) + \text{CO}_3^{2-}(\text{aq}) \rightarrow \text{CaCO}_3(\text{s})$	1 point is earned for the correct equation.
---	---

- (b) The diagram below is incomplete. Draw in the species needed to accurately represent the major ionic species remaining in the solution after the reaction has been completed.



The drawing shows one Ca^{2+} ion.	1 point is earned for drawing a Ca^{2+} ion.
---	---

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 3 (continued)

The student filters and dries the precipitate of CaCO_3 (molar mass 100.1 g/mol) and records the data in the table below.

Volume of Na_2CO_3 solution	50.0 mL
Volume of 1.0 M $\text{Ca}(\text{NO}_3)_2$ added	100.0 mL
Mass of CaCO_3 precipitate collected	0.93 g

(c) Determine the number of moles of Na_2CO_3 in the original 50.0 mL of solution.

$0.93 \text{ g CaCO}_3 \times \frac{1 \text{ mol CaCO}_3}{100.1 \text{ g}} = 0.0093 \text{ mol CaCO}_3$ $0.0093 \text{ mol CaCO}_3 \times \frac{1 \text{ mol Na}_2\text{CO}_3}{1 \text{ mol CaCO}_3} = 0.0093 \text{ mol Na}_2\text{CO}_3$	1 point is earned for the correct answer.
--	---


(d) The student realizes that the precipitate was not completely dried and claims that as a result, the calculated Na_2CO_3 molarity is too low. Do you agree with the student's claim? Justify your answer.

Disagree. The presence of water in the solid will cause the measured mass of the precipitate to be greater than the actual mass of CaCO_3 . As a result, the calculated number of moles of CaCO_3 and moles of Na_2CO_3 will be greater than the actual moles present. Therefore the calculated concentration of $\text{Na}_2\text{CO}_3(aq)$ will be too high.	1 point is earned for the correct answer with valid justification.
---	--

(e) After the precipitate forms and is filtered, the liquid that passed through the filter is tested to see if it can conduct electricity. What would be observed? Justify your answer.

The liquid conducts electricity because ions ($\text{Na}^+(aq)$, $\text{Ca}^{2+}(aq)$, and $\text{NO}_3^-(aq)$) are present in the solution.	1 point is earned for the correct answer with valid justification.
--	--

The student decides to determine the molarity of the same Na_2CO_3 solution using a second method. When Na_2CO_3 is dissolved in water, $\text{CO}_3^{2-}(aq)$ hydrolyzes to form $\text{HCO}_3^-(aq)$, as shown by the following equation.



AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 3 (continued)

(f) The student decides to first determine $[\text{OH}^-]$ in the solution, then use that result to calculate the initial concentration of $\text{CO}_3^{2-}(\text{aq})$.

(i) Identify a laboratory method (not titration) that the student could use to collect data to determine $[\text{OH}^-]$ in the solution.

Determine the pH of the solution using a pH meter.	1 point is earned for identifying a valid method.
--	---

(ii) Explain how the student could use the measured value in part (f)(i) to calculate the initial concentration of $\text{CO}_3^{2-}(\text{aq})$. (Do not do any numerical calculations.)

<p>First determine $[\text{OH}^-]$ using $\text{pOH} = 14 - \text{pH}$, then $[\text{OH}^-] = 10^{-\text{pOH}}$. Then, use the K_b expression and an ICE table (see example below) to determine $[\text{CO}_3^{2-}]$ and $[\text{HCO}_3^-]$ at equilibrium. The initial concentration of CO_3^{2-}, c_i, is equal to the sum of the equilibrium concentrations of CO_3^{2-} and HCO_3^-.</p> <table border="1" style="margin-left: auto; margin-right: auto; border-collapse: collapse; text-align: center;"> <thead> <tr> <th></th> <th>$\text{CO}_3^{2-}(\text{aq}) + \text{H}_2\text{O}(\text{l})$</th> <th>$\rightleftharpoons$</th> <th>$\text{HCO}_3^-(\text{aq}) + \text{OH}^-(\text{aq})$</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>c_i</td> <td>---</td> <td>0</td> </tr> <tr> <td>C</td> <td>$-x$</td> <td>---</td> <td>$+x$</td> </tr> <tr> <td>E</td> <td>$c_i - x$</td> <td>---</td> <td>x</td> </tr> </tbody> </table> $K_b = \frac{(x)(x)}{c_i - x} \Rightarrow c_i = \frac{(x)(x)}{K_b} + x$		$\text{CO}_3^{2-}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	\rightleftharpoons	$\text{HCO}_3^-(\text{aq}) + \text{OH}^-(\text{aq})$	I	c_i	---	0	C	$-x$	---	$+x$	E	$c_i - x$	---	x	<p>1 point is earned for a valid method of determining $[\text{OH}^-]$ from the measured value.</p> <p>1 point is earned for a valid method of determining the <u>initial</u> concentration of CO_3^{2-}.</p>
	$\text{CO}_3^{2-}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	\rightleftharpoons	$\text{HCO}_3^-(\text{aq}) + \text{OH}^-(\text{aq})$														
I	c_i	---	0														
C	$-x$	---	$+x$														
E	$c_i - x$	---	x														

(g) In the original Na_2CO_3 solution at equilibrium, is the concentration of $\text{HCO}_3^-(\text{aq})$ greater than, less than, or equal to the concentration of $\text{CO}_3^{2-}(\text{aq})$? Justify your answer.

Less than. The small value of K_b , 2.1×10^{-4} , indicates that the reactants are favored.	1 point is earned for the correct answer with a valid justification.
--	--

(h) The student needs to make a $\text{CO}_3^{2-}/\text{HCO}_3^-$ buffer. Is the Na_2CO_3 solution suitable for making a buffer with a pH of 6? Explain why or why not.

<p>No, the Na_2CO_3 solution is not suitable. The $\text{p}K_a$ of HCO_3^- is 10.32. Buffers are effective when the required pH is approximately equal to the $\text{p}K_a$ of the weak acid. An acid with a $\text{p}K_a$ of 10.32 is not appropriate to prepare a buffer with a pH of 6.</p>	<p>1 point is earned for the correct answer with a valid explanation.</p>
--	---