

**AP<sup>®</sup> CHEMISTRY**  
**2019 SCORING GUIDELINES**

**Question 4**

A student is doing experiments with  $\text{CO}_2(g)$ . Originally, a sample of the gas is in a rigid container at 299 K and 0.70 atm. The student increases the temperature of the  $\text{CO}_2(g)$  in the container to 425 K.

(a) Describe the effect of raising the temperature on the motion of the  $\text{CO}_2(g)$  molecules.

The average speed of the molecules increases as temperature increases.	1 point is earned for the correct answer.
--	---

(b) Calculate the pressure of the  $\text{CO}_2(g)$  in the container at 425 K.

Both the volume and the number of molecules are constant, therefore $\frac{P_1}{T_1} = \frac{P_2}{T_2} \Rightarrow \frac{0.70 \text{ atm}}{299 \text{ K}} = \frac{P_2}{425 \text{ K}} \Rightarrow P_2 = 0.99 \text{ atm}$	1 point is earned for the correct answer.
--	---

(c) In terms of kinetic molecular theory, briefly explain why the pressure of the  $\text{CO}_2(g)$  in the container changes as it is heated to 425 K.

Faster-moving gas particles collide more frequently with the walls of the container, thus increasing the pressure. OR Faster-moving gas particles collide more forcefully with the walls of the container, thus increasing the pressure.	1 point is earned for a correct explanation.
--	--

(d) The student measures the actual pressure of the  $\text{CO}_2(g)$  in the container at 425 K and observes that it is less than the pressure predicted by the ideal gas law. Explain this observation.

The attractive forces between $\text{CO}_2$ molecules result in a pressure that is lower than that predicted by the ideal gas law.	1 point is earned for a correct explanation.
--	--